

by the conductivity, using the word "conductivity" (as above explained) in such a sense as to include convection. The rate of increase downwards must, therefore, be the same at all depths at which this conductivity is the same.

This reasoning applies to superposed strata at the same place, and assumes them to be sufficiently regular in their arrangement to ensure that the flow of heat shall be in parallel lines, not in converging or diverging lines.

5. If we have reason to believe that the flow of heat upwards is nearly the same at all places, then the above reasoning can also be applied approximately to the comparison of one place with another—that is to say, the rates of increase downwards, in two masses of rock at two different places, must be approximately in the inverse ratio of their conductivities. In the cooling of a heated sphere of heterogeneous composition, the rates of flow would at first be very unequal through different parts of the surface, being most rapid through those portions of the substance which conducted best; but these portions would thus be more rapidly drained of their heat than the other portions, and thus their rates of flow would fall off more rapidly than the rates of flow in the other portions. If the only differences in the material were differences of conductivity, we might on this account expect the outflow to be after a long time nearly the same at all parts of the surface. But when we come to consider differences of "thermal capacity per unit volume," it is clear that with equal values of "diffusivity," that is of "conductivity divided by thermal capacity of unit volume" in two places, say in two adjacent sectors of the globe, there would be the same distribution of temperatures in both, but not the same flow of heat, this latter being greatest in the sector in which the capacity and conductivity were greatest.

Where we find, as in Mr. Deacon's observations at Bootle, near Liverpool, and to a less marked degree in the observations of Sir William Fairbairn and Mr. Garside, near Manchester, an exceptionally slow rate of increase, without exceptionally good conductivity, it is open to us to fall back on the explanation of exceptionally small thermal capacity per unit volume in the underlying region of the earth, perhaps at depths of from a few miles to a few hundred miles.

6. A question which was brought into consideration by Prof. Hull, in connection with the great difference between the rate of increase at Dukinfield and that at Rosebridge, is the effect of the dip of the strata upon the vertical conduction of heat. Laminated rocks conduct heat much better along the planes of lamination than at right angles to them. If k_1 denote the conductivity along, and k_2 the conductivity normal to the planes of lamination, and if these planes are inclined at an angle θ to the horizon, the number of feet per degree of increase downwards corresponding to a given rate of outflow through the surface, will be the same as if the flow were vertical with a vertical conductivity:—

$$k_1 \sin^2 \theta + k_2 \cos^2 \theta.$$

Prof. Herschel finds about 1·3 as the ratio of the two principal conductivities in Loch Rannoch flagstone, and 1·875 as the ratio in Festiniog slate.

The dip of the strata at Dukinfield is stated by Mr. Garside to be 15° , and we have $\sin^2 15^\circ = \cdot 07$, $\cos^2 15^\circ = \cdot 93$.

If we assume $k_1 = 1\cdot3 k_2$, as in the case of flagstone, we find for the effective vertical conductivity $k_2 (\cdot 09 + \cdot 93) = 1\cdot02 k_2$, so that the number of feet per degree would only be increased by 2 per cent.

It is not likely that the two conductivities in the strata at Dukinfield are so unequal as even in the case of flagstone, so that 2 per cent. is a high estimate of the effect of their dip on the vertical rate of increase so far as pure conduction is concerned. The effect of dip in promoting the percolation of water is a distinct consideration, but the workings of the Dukinfield mines are so dry that this action does not seem to be important.¹

(To be continued.)

METAMORPHIC ROCKS OF BERGEN²

THE metamorphic rocks of the Bergen Peninsula in Norway continue to attract the attention of Norwegian geologists, and we have before us, as an addition to the well-known works

¹ Though the workings are dry there is a large quantity of water in the superincumbent strata.

² Hans H. Reusch, "Silurifossiler og Pressede Konglomerater i Bergenskjiffene."—Universitetsprogram for første Halvaar (1883). Kristiania, 1882.

of Naumann, Leopold von Buch, Esmark, Keilhau, Kjerulf, and Hjörtal, a new elaborate and interesting work by M. Hans H. Reusch, which deals with the same subject. These rocks consist, as is known, of a variety of quartziferous talc-mica schists, diorite, clay-slates, conglomerates, and strongly-developed gneisses and granites. Various and very different opinions have been expressed as to the origin of these rocks. The researches of M. Reusch give a key to this question, as he has discovered in the clay-slates, which seem to constitute the upper part of these vertical strata, numerous fossils belonging to the lower part of the Upper Silurian formations, namely *Halysites catenularia* and *Cyathophyllum*, changed into white calcareous spar, a few tubular bodies (presumably *Syringophyllum organum*), some gasteropods (*Murchisonia* or *Subulites*?) some trilobites, as *Calymene*, also *Phacops* or *Dalmanites*, and some brachiopods. The presence of these fossils is the more interesting as the whole series of schists was often considered as of igneous origin. As to the gneisses and gneisso-granites of the peninsula, M. Reusch has given great attention to their structure and to the remarkable results of pressure which the rocks have undergone. He shows how granitic veins were folded and crumpled, how a kind of transversal stratification has arisen in beds of stratified gneiss under the influence of pressure, and he concludes, from an accurate study of the subject, that altogether the rocks show a far greater degree of plasticity than might have been supposed. "It seems that there are masses, as, for instance, the gneiss of Svenningdal, that have on one side a true stratified structure (not merely parallel or schistose structure) which could hardly be found in a rock of igneous origin, and on the other side send veins, or have included fragments which have undergone metamorphic changes."

One of the most attractive features of M. Reusch's work is the attention he has given to metamorphic phenomena and to changes caused by the pressure undergone by strata during their folding. The metamorphic phenomena were especially studied in the Osören district. The limestone which contains Silurian fossils has become marble, and the cause of metamorphism was not contact with some eruptive rock, but rather (as was observed in the Bernese Oberland by Swiss geologists) pressure and the molecular movements which pressure has occasioned in rocks. The clay, in which trilobites and other animals were entombed at Vagtdalen, has become a rock like muscovite-schist with porphyritically inclosed clusters of mica. As to the gneiss which appears among undoubtedly Silurian rocks, the author is inclined to consider it as sedimentary and as having been originally formed of loose materials. The granulite is clearly stratified and of sedimentary origin. The changes produced by the folding of strata and by the pressure they have undergone, are described with much accuracy and illustrated by many drawings. The fossils are nearly all compressed and elongated; the formerly conical coralla have received the shape of flat elongated biscuits, in accordance with the direction of pressure and stretching. The same is true with regard to all other fossils. In the green conglomerates at Osören, all the stones are flattened and elongated, acquiring thus a shape which they could not have possessed originally; very many of them have such a shape as to give in a cross-section the form of a lance-shaped leaf. The same structure, remarks the author, may be observed with the aid of a microscope in the "hones" from Eidsmarken in the South of Norway. Altogether the work of M. Reusch, although not rich in conclusions and generalisations, will be a welcome addition to the accurate knowledge of the still little understood metamorphic rocks. The Norwegian text of this work is accompanied with a rather too short *résumé* in English.

P. K.

CHEMICAL NOTES

CARNELLEY [*Chem. Soc. Jnl.* Trans., 1881, p. 317] has repeated his experiments on the effect of pressure on the melting-point of mercuric chloride, and has obtained results which show that this salt cannot be obtained in the solid state at temperatures appreciably above its melting-point.

JAHN [*Berliner Berichte*, xv. p. 1238] has made a series of careful determinations of the density of bromine vapour, which, when compared with similar observations on chlorine made by Ludwig, show that bromine vapour does not attain the normal density ($\text{Br} = 79\cdot95$) until it is heated to 160° above its boiling-point; and also that although chlorine exhibits smaller divergences from the normal density than bromine, it nevertheless

must be heated about 240° above its boiling-point before its density becomes strictly normal. From the experiments of V. Meyer and others, iodine vapour appears to be normal a very few degrees above the boiling-point. It appears, therefore, probable that vapour of chlorine, bromine, or iodine at low temperatures, contains groups of molecules which are dissociated as the temperature rises, and that the greater the molecular weight the more easily are these molecular groups dissociated.

REFERENCE was made in a note (NATURE, vol. xxvi. p. 306), to Spring's researches on the expansion of isomorphous compounds; in last number of the *Berliner Berichte* Pettersson draws attention to accurate determinations of the specific gravities of various alums, published by him a few years ago, which proved that the quotients of the specific gravities of the alums by their respective formula weigh's, are not equal, as assumed by Spring, but show considerable differences. Spring's work on the expansion of alums may lead to interesting results, but it seems evident that he has been too hasty in drawing sweeping conclusions regarding the molecular structure of solids from quite insufficient data.

HERR G. KRUSZ describes [*Berliner Berichte*, xv. 1243] a spectroscopic method for determining whether there is, or is not, any chemical action occurring in a solution containing two or more coloured salts. The method consists essentially in comparing the sums of the absorption spectra of the individual salts with the absorption spectrum of the solution containing all the salts.

AN abstract of an important paper by Mendeleeff on thermochemistry appeared in the *Berichte* for July 10 [xx. 1555]. Mendeleeff asserts that the data hitherto attained by Berthelot, Thomsen, and others, regarding the "heats of formation" of hydrocarbons stand in need of correction, because allowance has not been made for the physical changes, involving absorption or evolution of heat, which in every case accompany the chemical changes considered. Mendeleeff gives a table showing the "heats of formation" from marsh gas, carbon monoxide, and carbon dioxide, of a series of hydrocarbons; the chemical reactions, the thermal equivalents of which are set down in this table, are reactions which actually occur, unlike the reactions of formation of Berthelot and others, which as a rule cannot be realised in actual experiments.

MESSRS. SMITH AND LOWE find that when chlorine is passed through a porcelain tube heated to 1030° , and then into potassium iodide solution, less iodine is liberated than is the case when the same quantity of chlorine is allowed to act on the iodide at ordinary temperature; they conclude, therefore, that chlorine is partly dissociated at a temperature of 1030° (*Chem. News*, xlv. 226).

ACCORDING to Mixer (*Amer. Chem. J.*, iv. 35), urea is readily obtained by passing ammonia and carbon dioxide through a red-hot tube; ammonium cyanate is probably produced, and then transformed into urea.

SELMI (*R. Acad. dei Lincei*, v. 174) states that he has found alkaloidal compounds having specific poisonous actions in the urine of patients suffering from paralysis, tetanus, &c. He considers death to be determined by the action of these poisons produced by the progress of the disease.

By electrolysis of water by a powerful current, using a positive electrode of gas-coke purified by the action of chlorine at a very high temperature, Bartoli and Papasogli (*Gazzetta Chim. Ital.*, 1882, 113) obtained a black solution, which, when acidified with hydrochloric acid, yielded a black substance having the composition $C_{11}H_2O_4$ (when dried at 140°). The properties of this substance—*Mellogen*—are very peculiar; in some points it resembles graphitic acid; it dissolves in water to form an inky-black neutral liquid; on exposure to air or by the action of oxidising agents it yields mellitic acid and other acids, which are generally regarded as addition products of benzene.

UNIVERSITY AND EDUCATIONAL INTELLIGENCE

PROF. BONNEY begins his course of lectures on Petrology, Physiography, and Stratigraphical Geology at University College, Gower Street, on October 10, at twelve o'clock. The course will extend over two terms. Classes will also be formed for catechetical instruction and for the study of the microscopic structure of rocks.

SOCIETIES AND ACADEMIES

PARIS

Academy of Sciences, September 25.—M. Blanchard in the chair.—A report was given of the ceremony at the recent inauguration of a statue to Antoine-César Becquerel at Châtillon-sur-Loing, on September 24, when addresses were delivered by MM. Cochéry, Dumas, Fremy, Mercadier, Barral, and the Mayor.—On a question of principle which relates to the theory of shock of imperfectly elastic bodies, by M. Resal.—Outbreaks of plague in Kurdistan during the last twelve years, by M. Thälözan. The facts are thought to afford further proof of the independence of most of the plague-centres, the small tendency of the disease to spread beyond a small number of localities, and the limited duration of the epidemics, even in their gravest form. The north and north-west of Persia are the parts where plague-epidemics are less rare.—Possibility of introducing a tube into the larynx without producing pain or any reaction, by M. Brown-Séquard. He produces local anaesthesia in mammalia by directing a rapid current of carbonic acid on the upper part of the larynx (through an incision), for a variable time (fifteen seconds to two or three minutes). The effect lasts two to eight minutes after stoppage.—A telegram from the Emperor of Brazil (dated Rio, September 12, 6h. 10m.), announced the observation (at Rio Observatory) of a brilliant comet; estimated position in the morning, ascension 10h., declination 2° S. "probably Pona's comet expected.—On a comet observed at Nice, by MM. Thollon and Gouy. This was visible near the sun on September 18, at mid-day, to the naked eye, when the direct solar rays were masked. The spectrum had for essential character the presence of the bright lines of sodium (fine and perfectly separate) in the nucleus and parts near. A slight displacement was held to indicate withdrawal from the earth. No part of the comet showed bands of carbon, nor any band or line but those of sodium (probably because of a masking by diffuse light). On the morning of the 21st the comet had become invisible.—M. Flammarion communicated telegrams from Spain, Portugal, the South of France, Algeria, and Italy, announcing observations of a comet on September 17, 18, and 19.—On an observation of the great comet of 1882, seen from a balloon, by M. de Fonvielle. M. Mallet made the ascent at his request (having keener vision), and took some measurements. The diameter of the comet was about two-tenths of that of the sun, and the distance of the comet's centre from that of the sun about $2\frac{1}{3}$ subjective diameters of the sun. The cometary sphere was penetrated by an isosceles cone, symmetrically placed to the line of centres, penetrating to two-thirds of its vertical meridian plane. The length of the apothem of the cone was about a solar radius.—Description of a complete regular dodecahedron, by M. Barbier.—On the development of Alcyonarians, by MM. Kowalevsky and Marion.—On the histological structure of the digestive tube of *Holothuria tubulosa*, by M. Sourdau.—Analysis of the milk of Galibi women at the Jardin d'Acclimatation, by Mme. Brés. The milk is rich in butter and lactose, and there is very little casein.

CONTENTS

PAGE

FLUIDS	545
HANDBOOK OF INVERTEBRATE ZOOLOGY. By Prof. E. RAY LANKESTER, F.R.S.	548
OUR BOOK SHELF:—	
White's "Cameos from the Silver-land"	548
LETTERS TO THE EDITOR:—	
The Recent Aurora.—Prof. J. P. O'REILLY: SYDNEY EVERSHED; W. J. LINEHAM; Dr. GUSTAVE PLARR; ELIZTH BROWN	548
An Insect attacking a Worm.—EDWIN LAWRENCE	549
White Ants.—Surgeon-Major C. BIDIE	549
BRITISH SPONGES	550
EXPLORATION IN SIAM	550
THE COMET	551
SPECTROSCOPIC WEATHER DISCUSSIONS. By Prof. C. PIAZZI SMYTH, Astronomer Royal for Scotland (With Diagrams)	551
ELECTRIC NAVIGATION. By Prof. SILVANUS P. THOMPSON (With Illustrations)	554
NOTES	556
BIOLOGICAL NOTES:—	
On a New Genus of Cryptophyceæ	557
Seed Leaves of <i>Bursera</i> (With Illustration)	558
Affinities of the Bower Birds	558
GEOGRAPHICAL NOTES	559
PELAGIC LIFE. By Prof. H. N. MOSELEY, F.R.S.	559
UNDERGROUND TEMPERATURE	564
METAMORPHIC ROCKS OF BERGEN	567
CHEMICAL NOTES	567
UNIVERSITY AND EDUCATIONAL INTELLIGENCE	568
SOCIETIES AND ACADEMIES	568